

# Technical Information Sheet



Envy is for use on three key areas: Where horses or cattle graze low input grassland
Replace straight fluroxypyr on chickweed
Weed control in new sown leys

# **Key facts**

| Product Registration Number: | PCS 05806  |  |  |  |
|------------------------------|--|--|--|--|
| Active Ingredient:           | 100 g / litre fluroxypyr + 2.5 g / litre florasulam  |  |  |  |
| Pack size:                   | 3.0 L  |  |  |  |
| Formulation:                 | Suspension emulsion  |  |  |  |
| Maximum Individual Dose:     | 2.0 L/ha to established grassland<br>1.5 L/ha to new sown leys and grass for seed  |  |  |  |
| Maximum Total Dose:          | 2.0 L/ha to established grassland<br>1.5 L/ha to new sown leys and grass for seed  |  |  |  |
| Application Timing:          | 1st February to 30th November for established grassland and new sown leys<br>1st March to 30th November for grass grown for seed |  |  |  |
| Water Volumes:               | 200 L/ha new sown leys, 200-400 L/ha established grassland   |  |  |  |
| Spray Quality:               | Medium as defined by BCPC  |  |  |  |

### Key benefits

- Perfect for paddocks:
  - Control buttercups, dandelions, daisies, plantains and other perennial weeds typically found in paddocks.
  - No manure residue restrictions.
  - Short stock withdrawal period of just 7 days so quick return of animals. Use at 2.0 L/ha.
- Controls chickweed up to flowering in established grassland. Better than straight fluroxypyr in fluctuating diurnal temperatures
- Valuable for use in new sown leys due to wide application window and broad spectrum of weeds that can be controlled. (Apply any time between 1st February and 30th November). Use at 1.5 L/ha.
- Ideal for use in grass seed crops
- 3 litre pack treats 1.5 ha (3.7 acres) at 2.0 L/ha rate, or 2 ha (4.9 acres) at 1.5 L/ha rate
- · Rainfast in 2 hours

## Best use advice

- In new sown leys apply to weeds which are small and actively growing, and when the grass has reached the 3 leaf stage.
- Envy is a contact herbicide so broad-leaved weeds not present at application will not be controlled
- · If weeds grow beyond the ideal stage, consider cutting the sward and spraying the regrowth
  - Allow 2-3 weeks after cutting before applications of Envy are made to allow sufficient regrowth
- · Allow an interval of 4 weeks after application before cutting grass to optimise the effect of the herbicide
- Do not roll or harrow grass for 10 days before or 7 days after application
- Do not spray where clover or other legumes are a valuable constituent of the sward
- Grass seed can be stitched in 1 month after application
- Clover can be stitched in 3 months after application
- Grazing can resume just 7 days after application
  - This may need to be longer if foliage of poisonous weeds such as ragwort is present
  - · Envy will kill clover
- To avoid subsequent injury to crops other than grassland and cereals, all spraying equipment must be thoroughly cleaned using All Clear Extra spray cleaner

Visit us at corteva.ie



# Weed Spectrum

Where we have knowledge of how our products might affect other grassland weeds we have detailed it in the following tables. These are not recommendations; just an indication of what effects might be achieved. • indicates information based on anecdotal or limited data, and as such the user bears the risk in respect of failures concerning the efficacy and phytotoxicity.

| Control of Annual Weeds with Envy 1.5 L    |                                    |                                   |              |                       |                                    |  |  |
|--|------------------------------------|-----------------------------------|--------------|-----------------------|------------------------------------|--|--|
| Black bindweed                             | ▲4TL                               | Fool's parsley                    | ▲4TL         | Рорру                 | 4TL                                |  |  |
| Black nightshade                           | ▲4TL                               | Forget-me-not                     | 4TL          | Redshank              | <b>≜</b> 2TL                       |  |  |
| Bristly ox-tongue                          | -                                  | Fumitory                          | <b>≜</b> 2TL | Scarlet pimpernel     | ▲2TL                               |  |  |
| Charlock                                   | 200mm                              | Groundsel                         | ▲2TL         | Shepherd's-<br>purse  | 4TL                                |  |  |
| Chickweed                                  | flowering                          | Hemp-nettle                       | ▲4TL         | Small nettle          | ▲2TL                               |  |  |
| Cleavers                                   | flowering                          | Himalayan<br>balsam               | -            | Speedwells            | <b> ¢</b> cotyledons               |  |  |
| Corn chamomile                             | 150mm                              | Knotgrass                         | 4TL          | Spurrey               | <b>≜</b> 2TL                       |  |  |
| Corn marigold                              | <b>▲</b> 150mm                     | Mayweed                           | 200mm        | Wild radish           | ▲2TL                               |  |  |
| Cranesbill                                 | ×                                  | Medick                            | -            | Yellow rattle         | -                                  |  |  |
| Dead-nettles                               | ⁴cotyledons                        | Orache                            | ×            |                       |                                    |  |  |
| Fat-hen                                    | ▲2TL                               | Pale persicaria                   | <b>≜</b> 2TL |                       |                                    |  |  |
| Control of Perennial Weeds with Envy 2.0 L |                                    |                                   |              |                       |                                    |  |  |
| Bramble                                    | <b>▲</b> ✓ ✓                       | Giant hogweed                     | ×            | Ox-eye daisy          | ×                                  |  |  |
| Broom                                      | ×                                  | Gorse                             | ×            | Plantain<br>(greater) | <b>A</b> $\checkmark$ $\checkmark$ |  |  |
| Burdock                                    | -                                  | Ground elder                      | ×            | Ragwort               | ×                                  |  |  |
| Buttercup                                  | <b>/ / /</b>                       | Ground ivy                        | -            | Ribwort<br>plantain   | <b>A</b> 🗸 🗸                       |  |  |
| Cinquefoil                                 | -                                  | Hawthorn                          | ×            | Rosebay<br>willowherb | -                                  |  |  |
| Clover, trefoil                            | <b>A</b> $\checkmark$ $\checkmark$ | Hemlock                           | <b>▲</b> √   | Rushes                | ×                                  |  |  |
| Coltsfoot                                  | -                                  | Horsetail<br>( <i>Equisetum</i> ) | ▲✓           | Self-heal             | -                                  |  |  |
| Common sorrel                              | <b>A</b> $\checkmark$ $\checkmark$ | Japanese<br>knotweed              | ×            | Silverweed            | -                                  |  |  |
| Cow parsley                                | ×                                  | Lesser<br>celandine               | -            | Thistles              | ×                                  |  |  |
| Daisy                                      | <b>√√√</b>                         | Mallow                            | <b>▲</b> √√  | Vetch, tare           | <b>▲</b> √√                        |  |  |
| Dandelion                                  | <b>/ / /</b>                       | Mugwort                           | -            | Yarrow                | * < < <                            |  |  |
| Docks                                      | <b>√</b> √                         | Nettles                           | <b>▲</b> √   | Yellow/Flag Iris      | -                                  |  |  |
| Field bindweed                             | ▲4TL                               | Old man's<br>beard                | -            |                       |                                    |  |  |

 $\checkmark$   $\checkmark$  ✓, Good control,  $\checkmark$  ✓ Moderate control,  $\checkmark$  Some control,  $\times$  No control, -No Information, TL True Leaves,  $^{\blacktriangle}$ Anecdotal or limited information

